



Agromet Advisory Bulletin for Thrissur District (Valid from 12.04.2023 to 16.04.2023)



Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 15/2023, (29) Tuesday

dated 11.04.2023

A. Significant weather for the preceding week at vellanikkara								
Rainfall	Maximum	Minimum	RH	RH	Wind speed	Sun shine	Evap. (mm)	
(mm)	Temp.(°C)	Temp.(°C)	FN (%)	AN (%)	(Km/h)	hours		
(0.0)	(34.4-38.3)	(24.0-25.9)	(79-89)	(31-64)	(2.2-3.0)	(7.5 -10.1)	(4.8-5.9)	

B. Medium range weather forecast of IMD for next five days						
Weather parameters	12.04.2023	13.04.2023	14.04.2023	15.04.2023	16.04.2023	
Rainfall (mm)	0.5	0	0	0	0.6	
Maximum temperature (°C)	38	38	38	38	36	
Minimum temperature (°C)	26	26	26	26	23	
Maximum Relative Humidity (%)	80	80	80	80	80	
Minimum Relative Humidity (%)	57	57	57	57	57	
Wind speed (kmph)	3	4	3	3	3	
Wind direction (deg)	250	230	270	250	270	
Total cloud cover (octa)	8	5	8	8	8	

Weather summary / Alert: Light rainfall is expected in isolated places of the district. Normal rainfall is expected in Kerala.

C. General and Agrometeorological Advisories for Thrissur District

The following are immediate steps to be followed to protect farmers, plants, and birds from the summer heat. Drink plenty of clean water from time to time to prevent dehydration. Avoid going out in hot weather. Thus you can get rid of sunburn. While working in the farm, try to use a hat or an umbrella to prevent direct sunlight. As the climate is hot and dry, irrigate the crops as required to ensure adequate soil moisture. Irrigate only, when there is no strong sunlight. Drip irrigation can reduce water loss to some extent.

Crop	Pest/ Disease	Crop/ Animal Advisory				
Puncha rice		Ensure water availability in paddy fields.				
(Milky stage)	1	This is especially important in late sowing fields				
Banana	Mites Since dry weather condition is prevailed, there is a chance of mite attack in banana. Apply wettable sulphur 3 gram per one litre of water. Give strong propping to banana.					
Brinjal	White fly	During high temperature, there is a chance of white fly attack in brinjal. Apply 2% neem oil garlic emulsion or spray 20g <i>Lecanicillium lecanii</i> per one litre of water in ten days intervals.				

Animal Husbandry:- During cloudy condition, avoid bathing the cow and washing the cattle shed, otherwise more heat will be build up. Provide proper air circulation using fans. Assure increased ventilation in the cattle shed. Ensure sufficient supply of clean drinking water to the animals. (CAADECCS, KVASU, Mannuthy).