



Agromet Advisory Bulletin for Thrissur District (Valid from 07.5.2022 to 11.5.2022)



Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 18/2022, (36) Friday

dated 6.5.2022

A. Significant weather for the preceding week at vellanikkara							
Rainfall	Maximum	Minimum	RH	RH	Wind speed	Sun shine	Evap. (mm)
(mm)	Temp.(°C)	Temp.(°C)	FN (%)	AN (%)	(Km/h)	hours	
(26.9)	(33.2-35.2)	(22.4-26.4)	(91-94)	(57-68)	(1.9-2.7)	(1.8-7.6)	(3.2-4.2)

B. Medium range weather forecast of IMD for next five days					
Weather parameters	07.5.2022	08.5.2022	09.5.2022	10.5.2022	11.5.2022
Rainfall (mm)	5	0	0	5	0
Maximum temperature (°C)	33	34	34	34	34
Minimum temperature (°C)	23	26	26	26	26
Maximum Relative Humidity (%)	89	89	89	89	89
Minimum Relative Humidity (%)	67	67	67	67	67
Wind speed (kmph)	3	3	3	3	3
Wind direction (deg)	250	250	250	340	320
Total cloud cover (octa)	7	4	4	7	3

Weather summary / **Alert:** Light rainfall is expected in the district and moderate rainfall is expected in isolated places. Below normal rainfall is expected in Kerala.

RAINFALL FORECAST FOR NEXT FIVE DAYS					
2022 May 07	2022 May 08	2022 May 09	2022 May 10	2022 May 11	
light rain	No rain	No rain	light rain	No rain	

C. General and Agrometeorological Advisories for Thrissur District

Ensure drainage facilities in low land. Proper mulching can be followed to conserve soil moisture.

Ensure dramage racinities in low land. I toper matering can be followed to conserve soil moistage.					
Crop	Pest/ Disease		Crop/ Animal Advisory		
Rice (virippu)	Land preparation	It is recommended to sow dhaincha seeds in the paddy field before virip planting			
Coconut	Management	Pits can be taken for planting new coconut seedlings.			
Banana	Sigatoka disease Quick wilt	As a p trichode	s a chance of sigatoka disease in banana. As a precaution, Spray 20 g monas per one liter of water. rophylactic measure to manage quick wilt in pepper, add 2 kg erma to a mixture of 90 kg FYM and 10 kg neem cake and kept for eks with sufficient moisture. Add 2.5 kg of this mixture to each		
Vegetables	Management	matured vine to control quick wilt of pepper. Nurseries can be prepared for vegetables.			
Animal	Care should be taken to prevent rat breeding in the premises of cattle shed to prevent				
Husbandry	leptospyrosis. (CAADECCS, KVASU, Mannuthy).				