




Agromet Advisory Service bulletin19/16, Friday dt.06-05-2016

(Issued jointly by IAAS Vellanikkara)

Dept. of Agriculture Meteorology, College of Horticulture, Vellanikkara

E mail: kauagmet@yahoo.co.in, cohagmet@kau.in



THRISSUR

CURRENT SYNOPTIC SITUATION:

An upper air trough extending up to 1.5 km above sea level runs from North Interior Karnataka to Comorin area

NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST

ISSUED ON: 06.05.2016

VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE : KERALA

PARAMETERS

ENSEMBLE FCST

	DAY-1 07/05	DAY-2 08/05	DAY-3 09/05	DAY-4 10/05	DAY-5 11/05
DISTRICT : THRISSUR					
Rainfall (mm)	1	18	6	16	2
Max Temperature (deg C)	36	35	35	35	35
Min Temperature (deg C)	25	25	24	24	24
Total cloud cover (octa)	6	6	6	6	6
Max Relative Humidity (%)	80	80	80	80	80
Min Relative Humidity (%)	60	60	60	60	60
Wind speed (kmph)	006	006	006	006	006
Wind direction (deg)	250	250	270	270	270

Agro Meteorological Advisories

Crop	Stage of the crop / Pest/Disease	Advisories
Paddy	Aphid	There is a chance of aphid attack in paddy. Spray Imidacloprid 3 milli in 10 litre of water.
Coconut	Leaf rot	Due to leaf rot the tender leaves of coconut decayed and dried of due to fungus, and the dried part get blown off in wind. This symptom seen in root(wilt) also. Spray bordeaux mixture in 5g per liter on the top of the plant. Proper irrigation and ploughing may help to improve the plant from the disease.
Banana	Leaf eating caterpillar	Spray 2ml of Quinalphos in one liter of water.
Vegetables (Amaranthus)	Leaf spot disease	Cut and remove the severely affected leaves. Apply 15 gm of Pseudomonas in cleared cow dung water after harvesting. Also apply Trichoderma mixed with cowdung in soil.