




# Agromet Advisory Service bulletin 20/17, Tuesday dt 16.05.2017

(Issued jointly by IAAS Vellanikkara)

Dept. of Agriculture Meteorology, College of Horticulture, Vellanikkara

E mail: kauagmet@yahoo.co.in, cohagmet@kau.in



## THRISSUR

### Significant weather for the preceding week at Vellanikkara

Rainfall weekly total	14.2mm
Maximum temperature	35.5-35.9 <sup>0</sup> C
Minimum temperature	23.7-26.4 <sup>0</sup> C
Max. RH	79-96%
Min. RH	45-66%
Wind speed	1.1-2.3 km/hr
Sun shine hour	0.1-9.5 hr
Evaporation	2.3-4.1 mm

**CURRENT SYNOPTIC SITUATION:** An upper air cyclonic circulation lies over Gulf of Martaban & extends upto 3.6km above sea level. An upper air east-west trough runs roughly along Lat 8<sup>0</sup>N between 2.1 & 3.6 km above sea level.

### NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST

ISSUED ON: 16.05.2017

VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE : KERALA

PARAMETERS	ENSEMBLE FCST				
	DAY-1 17/05	DAY-2 18/05	DAY-3 19/05	DAY-4 20/05	DAY-5 21/05
DISTRICT : THRISSUR					
Rainfall (mm)	0	5	15	0	2
Max Temperature ( deg C)	34	34	33	33	33
Min Temperature ( deg C)	26	26	25	25	25
Total cloud cover (octa)	4	6	8	4	5
Max Relative Humidity (%)	85	85	85	85	85
Min Relative Humidity (%)	60	60	60	60	60
Wind speed (kmph)	008	008	008	010	010
Wind direction (deg)	250	250	250	250	250

### Agro Meteorological Advisories:

Crop	Stage of the crop / Pest/Disease	Advisories
Banana	Sigatoka	Due to the changing weather pattern, there is a chance of sigatoka disease in banana. As a precaution, Spray Pseudomonas 20g per 1 liter of water. In case of severe attack spray 1 gm Bavistin by mixing it with gum in 1 litre of water on lower surface of leaf.
Pepper	Quick wilt	There is a chance of Quick wilt disease due to high relative humidity. Apply trichoderma or pseudomonas culture in soil or 1% Bordeaux mixture can be sprayed on the leaves and small branches.
Vegetables	Mosaic/ Yellowing	Due to the increase in atmospheric temperature, the attack of insects such as aphids, white flies, hoppers are more. As a result of this, there is a chance of mosaic disease in vegetables, which can be controlled by applying 5 g of Verticillium or Metarhizium in 1 litre of water. Add neem cake as manure. Neem based insecticides can be applied. Control the vectors by spraying Dimethoate 0.05 %. Uprooting and destruction of affected plants and collateral hosts should be done. In severe condition of mosaic, apply Thiomethoxan 0.2 g /L is effective. To control the vectors of yellowing spray 3ml nimbecidine or 1.5 ml Dimethoate in 1 litre of water.

**Preparation of Trichoderma culture:** Dry neem cake and powdered cowdung are mixed at 1:10 ratio and add 1 kg trichoderma with 100 kg of above mixture. Then sprinkle water and mix it thoroughly. Cover it with a perforated polythene sheet and keep it in shade for 5 days for multiplication. Again mix well and keep for 3 more days for further multiplication and incorporate into the soil for controlling Quick wilt.