



Agromet Advisory Bulletin for Thrissur District





AAS Bulletin No: 21/2023, (42) Friday

dated 26.05.2023

	Obse			ring last v to 26 th Ma			Weather Forecast (Valid for 27 th to 31 st May 2023)			y			
20/05	21/05	22/05	23/05	24/05	25/05	26/05	Date		27/05	28/05	29/05	30/05	31/05
0.0	0.2	0.8	0.5	0.0	1.5	5.6	Rainfall (mm)		5.0	6.0	0.1	4.0	7.0
34.3	34.4	34.9	35.0	34.8	34.2	34.8	$T_{max}(^{0}C)$		35	35	35	36	36
26.7	27.1	23.7	25.0	25.7	25.2	24.2	T _{min} (⁰ C)		24	25	26	26	27
Cloudy	Cloudy	Partially Cloudy	Cloudy	Partially Cloudy	Cloudy	Overcast	AM	Cloud	Overcast	Overcast	Cloudy	Overcast	Cloudy
Cloudy	Partially Cloudy	Overcast	Cloudy	Cloudy	Partially Cloudy		PM	cover					
91	95	89	84	86	92	93	RH-I (%)		85	90	93	92	91
62	59	58	59	63	57	77	RH-II (%)		66	58	64	64	59
2.3	2.2	2.5	2.1	1.9	2.3	2.1	Wind Speed (km/hr.)		8	10	8	8	8
С	С	С	С	С	С	С	AM	Wind	NW	NW	NW	NW	NW
SW	NW	ENE	SSW	WSW	NW	WNW	PM	direction					
Ra	Rainfall (mm) in last week Rainfall (mm)				Rainfall (mm) from 01/01/2023 to till dated				Rainfall (mm) from 01/03/2023 to today				
	8.6				133.2					133.2			

Weather summary / Alert: As per the forecast of IMD, there is a chance of light rainfall in the district and moderate rainfall at isolated places. Normal rainfall is expected in Kerala.

C. General and Agrometeorological Advisories for Thrissur District

Special crop management is required for agricultural crops before monsoon season. Provide supports for the bananas. Make the vegetable pandals stronger. Avoid standing in open areas to protect farmers and livestock from lightning and thunder. Proper mulching can be followed to conserve soil moisture.

Crop	Pest/ Disease	Crop/ Animal Advisory					
Rice (virippu)	-	Land preparation and nursery for virippu paddy (Rainfed) can be started after summer rains.					
Coconut & Arecanut	-	Pits can be taken for planting seedlings of coconut and arecanut.					
Banana	Sigatoka disease	There is a chance of sigatoka disease in banana. As a prophylactic measure to sigatoka leaf spot disease, spray 1% Bordeaux mixture. If Sigatoka attack is seen, spray one milli litre hexaconazole or one milli litre propiconazole per one liter of water on both the surface of the leaf.					

Animal Husbandry:- Care should be taken to avoid fungal growth in the cattle feed. During cloudy condition, avoid bathing the cow and washing the cattle shed, otherwise more heat will be build up. Provide proper air circulation using fans. Assure increased ventilation in the cattle shed. (CAADECCS, KVASU, Mannuthy).