




**Agromet Advisory Service bulletin 25/16, Tuesday dt 14-06-16**  
 (Issued jointly by IAAS Vellanikkara)  
 Dept. of Agriculture Meteorology, College of Horticulture, Vellanikkara  
 E mail: kauagmet@yahoo.co.in, cohagmet@kau.in  
**THRISSUR**



**Significant weather for the preceding week at Vellanikkara**

Rainfall weekly total	190.5 mm
Maximum temperature	26.6-31.8 <sup>0</sup> C
Minimum temperature	21.1-23.2 <sup>0</sup> C
Max. RH	95-98%
Min. RH	67-98 %
Wind speed	0.5-1.9 km/hr
Sun shine hour	0.0-3.0 hr
Evaporation	1.8-3.1 mm

**CURRENT SYNOPTIC SITUATION:**

A feeble trough and low pressure on sea level chart lies off Karnataka –Kerala coast .

**NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST**  
 ISSUED ON: 14.06.2016  
 VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE : KERALA

PARAMETERS	ENSEMBLE FCST				
	DAY-1 15/06	DAY-2 16/06	DAY-3 17/06	DAY-4 18/06	DAY-5 19/06
DISTRICT : THRISSUR					
Rainfall (mm)	5	07	22	40	40
Max Temperature ( deg C)	31	31	31	31	31
Min Temperature ( deg C)	22	22	22	22	22
Total cloud cover (octa)	5	5	6	8	8
Max Relative Humidity (%)	90	90	90	95	95
Min Relative Humidity (%)	80	80	80	80	80
Wind speed (kmph)	008	008	008	008	008
Wind direction (deg)	290	290	290	270	270

**Agro Meteorological Advisories:**

Crop	Stage of the crop / Pest/Disease	Advisories
Paddy (Jyothi, Kanchana)		Transplanting stage in first paddy crop.
Coconut	Bud rot disease	Cut and remove all the affected tissues of the crown. Apply Bordeaux paste and protect it from rain till normal shoot emergence. Transplanting of seedlings can be done during the week.
Banana	Pseudostem Weevil	Apply 2% neem oil-garlic-soap emulsion in the starting stage. In case of severe attack apply 3ml Chlorpyrifos in 1 liter of water, mix with gum, on the pseudostem as a paste.
Vegetables	Cow pea	This is the suitable time for cowpea planting.