



## **Agromet Advisory Bulletin for Thrissur District**





AAS Bulletin No: 32/2024, (63) Tuesday

Dated 06.08.2024

Observe		r during la	×	Dated 31s	<sup>t</sup> July		Weather Parameters		Weather Forecast Valid for 07 <sup>th</sup> August to 11 <sup>th</sup> August 2024				
31/07	01/08	02/08	03/08	04/08	05/08	06/08	Date		07/08	08/08	09/08	10/08	11/08
32.1	25.1	17.2	1.2	12.1	0.5	22.2	Rainfall (mm)		5	4	0.1	2	10
25.9	28.4	28.5	30.6	31.0	32.5	31.9	$T_{max}(^{0}C)$		32	32	32	31	31
24.0	23.5	23.9	22.9	23.6	24.1	24.5	$T_{min}(^{0}C)$		25	25	24	24	24
Overcast	Cloudy	Overcast	Partly Cloudy	Overcast	Cloudy	Overcast	AM	Cloud	Overcast	Overcast	Cloudy	Cloudy	Overcast
Overcast	Cloudy	Cloudy	Cloudy	Cloudy	Cloudy	96	PM	cover					
98	98	96	93	99	96		RH-	(%)	90	90	90	90	90
86	92	78	77	67	66		RH-II (%)		70	70	70	70	70
0.7	1.4	1.0	1.7	1.3	2.1	1.9	Wind Speed (km/hr.)		5	14	8	9	7
C	C	C	C	C	C	NNW	AM	Wind	WNW	NNW	NNW	WNW	WNW
WSW	C	SW	NNE	W	E		PM	direction					
Ra	Rainfall (mm) in last week Rainfall (mm) from 0						/01/2024 to till dated Rainfall (mm) from 01/06/2024 to today				o today		
110.4				2322.8					1692.0				

Weather summary / Alert: As per the forecast of IMD, there is a chance of moderate rainfall in isolated places of the district. Above normal rainfall is expected in Kerala.

## General and Agrometeorological Advisories for Thrissur District

Special crop management is required for agricultural crops during monsoon season. Proper drainage facilities have to be ensured. If the soil is more acidic, add lime as per soil testing. Pesticides should not be applied during rainy period. Provide supports to banana crops. Make the vegetable pandals stronger. To protect farmers and livestock from lightning and thunder avoid standing in open areas.

$\mathcal{O}$		$\mathcal{E}$					
Crop		Crop/ Animal Advisory					
Rice (virippu) Tillering stage	Rice case worm	During rainy season, there is a chance of rice case worm in paddy field. Drain water from the field for three days. To control this 25 kg sawdust mixed with one litre kerosene can be spread per one acre.					
Pepper	Quick wilt	Proper drainage facilities have to be ensured. As a prophylactic measure to control quick wilt of pepper, add 2 kg Trichoderma to a mixture of 90 kg FYM and 10 kg neem cake and kept for two weeks with sufficient moisture. Add 2.5 kg of this mixture to the base of each matured pepper vine.					
Cucurbitaceous Vegetables	Downy mildew	During rainy season, there is a chance of downy mildew attack in cucurbitaceous vegetables. Spray 2.5gm Mancozeb per one litre of water under the leaves.					
Nutmeg	Leaf fall	During rainy season, there is a chance of leaf fall in nutmeg. As a prophylactic measure to manage leaf fall in nutmeg, spray 1% Bordeaux mixture. Proper drainage facilities have to be ensured					
Animal Husbandry, Cattle food should be stored on a wooden board or iron board. If moisture is in the							

Animal Husbandry: Cattle feed should be stored on a wooden board or iron board. If moisture is in the cattle feed it may cause deadly fungal growth. Animals should not be allowed to go to water logged areas in order to avoid anthrax and foot lesions. (CAADECCS, KYASU, Mannuthy)