



Agromet Advisory Bulletin for Thrissur District (Valid from 12.08.2023 to 16.08.2023)



Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 32/2023, (64)Friday

Dated 11.08.2023

	Observed ated 5 th		_	st week gust 2023)			Weather	Parameters	Weather Forecast Valid for 12 th August to 16 th August 2023)				
05/08	06/08	07/08	08/08	09/08	10/08	11/08	Date		12/08	13/08	14/08	15/08	16/08
5.0	0.1	0.0	0.0	0.0	0.0	0.0	Rainfall (mm)		1	0.5	1	0.1	0.3
31.9	32.4	33.0	32.5	31.7	32.9	32.5	T max(⁰ C)		33	33	33	33	33
23.6	24.5	24.5	25.0	24.2	25.2	25.7	T min(⁰ C)		26	26	26	26	26
Partially Cloudy	Partially Cloudy	Cloudy	Cloudy	Cloudy	Cloudy	Cloudy	AM	Cloud	Overcast	Overcast	Overcast	Cloudy	Cloudy
Cloudy	Cloudy	Partially Cloudy	Cloudy	Cloudy	Cloudy		PM	cover					
96	95	92	93	89	92	93	RH-I (%)		96	92	92	92	92
63	61	61	66	62	62		RH-II (%)		78	72	72	72	72
1.8	1.7	2.0	1.9	1.5	1.7	1.6	Wind Speed (km/hr.)		4	4	4	4	3
С	С	С	С	С	С	С	AM	Wind	WNW	WNW	NW	NW	NNW
SSW	NW	WNW	NNW	WSW	NE		PM	direction					
Rainfall (mm) in last week				Rainf	Rainfall (mm) from 01/01/2023 to till dated				Rainfall (mm) from 01/06/2023 to today				
5.1				1098.3					964.9				

Weather summary / Alert: As per the forecast of IMD, intensity of rainfall is expected to decrease in the district. Below normal rainfall is expected in Kerala.

C. General and Agrometeorological Advisories for Thrissur District

Special crop management is required for agricultural crops during monsoon season. Proper drainage facilities have to be ensured. Propping of banana plants is recommended. Additional support for vegetable pandals should be strengthened. Pesticides should not be applied during rainy period.

Сгор	Pest/ Disease	Crop/ Animal Advisory						
Paddy (virippu) Tillering stage	In order to reduce the micronutrient deficiency in paddy, spray 10 grams of sampoorna mixture of paddy in one liter of water. Sampoorna mixture of paddy is available at the Kerala Agricultural University and it contains micro nutrients like magnesium, sulphur, boron, zinc, copper, iron, manganese and molybdenum.							
Coconut	25 to 50 kilogram of organic manure can be applied per one matured palm during rainy season by opening the coconut basin. Also, 0.5 kg of magnesium sulphate should be added in the months of August and September.							
Banana	It is good to add 500 grams of lime to banana plants at the time of planting in the months of August and September. Also, at the time of planting10 kg of green manure, compost or green leaves should be added or within a month after planting.							
Vegetables ((Brinjal, Tomatto, Chilly)	167 grams of urea and 84 grams of muriate of potash per cent can be applied after 20 to 30 days of planting brinjal, tomato and chilly.							
Animal Husbandry: - Precautions should be taken to avoid ectoparasitic infestations. Deworming of animals should be done after microscopical examination of dung sample in a veterinary hospital. Since there is a chance of mixing of contaminated water in water resources, apply disinfectants available in the market as per manufacturer's instructions. (CAADECCS, KVASU, Mannuthy).								