



## **Agromet Advisory Bulletin for Thrissur District**





Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 33/2022, (65) Tuesday

dated 16.8.2022

A. Significant weather for the preceding week at vellanikkara							
Rainfall	Maximum	Minimum	RH	RH	Wind speed	Sun shine	Evap. (mm)
(mm)	Temp.(°C)	Temp.(°C)	FN (%)	AN (%)	(Km/h)	hours	
(40.2)	(29.4-31.9)	(22.7-24.4)	(92-98)	(57-78)	(1.8-4.6)	(1.3-9.2)	(2.0-3.7)

B. Medium range weather forecast of IMD for next five days					
Weather parameters	17.08.2022	18.08.2022	19.08.2022	20.08.2022	21.08.2022
Rainfall (mm)	0.1	1	0.5	1	0.4
Maximum temperature (°C)	32	31	31	30	30
Minimum temperature (°C)	24	23	24	24	24
Maximum Relative Humidity (%)	92	92	92	92	92
Minimum Relative Humidity (%)	60	60	60	60	60
Wind speed (kmph)	6	8	6	10	8
Wind direction (deg)	270	270	270	270	290
Total cloud cover (octa)	5	6	6	6	6

Weather summary / Alert: Very light rainfall is expected in the district. Below normal rainfall is expected in Kerala.

RAINFALL FORECAST FOR NEXT FIVE DAYS					
2022 August 17	2022 August 18	2022 August 19	2022 August 20	2022 August 21	
Very Light rain	Very Light rain	Very Light rain	Very Light rain	Very Light rain	

## C. General and Agrometeorological Advisories for Thrissur District

Since rainfall is reduced, crop protection measures such as application of fertilizers and pesticides can be adopted for the crops. Copra and arecanut can be sun dried.

Crop	Pest/ Disease		Crop/ Animal Advisory		
Paddy (virippu) Tillering stage	Management  In order to reduce the micronutrient deficiency in paddy, spray 10 grasampoorna mixture of paddy in one liter of water. Sampoorna mixture of pavailable at the Kerala Agricultural University and it contains micro nutries magnesium, sulphur, boron, zinc, copper, iron, manganese and molybor Fertilizers like 20 kg of urea and 10 to 12 kg of potash per acre can be apply short duration and medium duration varieties.				
Banana	Rhizome rot	Rhizome rot Banana is prone to rhizome rot due to continuous rainfall. As a preventive measure, mix 20 grams of Pseudomonas in one liter of water and drench each banana basin with three liters of pseudomonas solution.			
Penner Foot rot continuous rainfall. As a prophylact		is a chance of foot rot disease in pepper due to high humidity caused by uous rainfall. As a prophylactic measure, spray 1% Bordeaux mixture. As a ation 2.5 to 3 grams of copper oxy chloride per one litre of water can be need at the basin for 3 to 5 times.			
Ginger	Stem borer		is a chance of stem borer in ginger. Spray 20 g of Beauveria per one litre of to control stem borer.		

Animal Husbandry- Deworming of animals should be done only after dung sample examination. In calves first dose of deworming medicine can be given within 15 days of age. (CAADECCS, KVASU, Mannuthy).