



Agromet Advisory Bulletin for Thrissur District

(Valid from 17.08.2024 to 21.08.2024)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara
based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 33/2024, (66) Friday

Dated 16.08.2024

Observed weather during last week (Dated 10 th August to 16 th August 2024)							Weather Parameters		Weather Forecast Valid for 17 th August to 21 st August 2024				
10/08	11/08	12/08	13/08	14/08	15/08	16/08	Date		17/08	18/08	19/08	20/08	21/08
26.1	1.4	2.8	0.0	39.8	0.2	0.0	Rainfall (mm)		7	6	8	8	14
31.7	31.0	31.7	32.7	34.5	30.5	31.4	T _{max} (⁰ C)		32	32	31	31	31
24.5	25.4	25.3	25.2	24.0	24.5	25.5	T _{min} (⁰ C)		25	25	24	24	24
Overcast	Cloudy	Cloudy	Cloudy	Cloudy	Cloudy	Overcast	AM	Cloud cover	Overcast	Cloudy	Overcast	Cloudy	Cloudy
Cloudy	Overcast	Cloudy	Cloudy	Overcast	Cloudy		PM						
98	95	97	89	95	93	93	RH-I (%)		90	90	90	88	88
73	70	70	75	80	76		RH-II (%)		76	76	76	72	72
2.4	1.7	2.2	2.0	1.9	2.2	1.4	Wind Speed (km/hr.)		4	4	4	3	4
C	C	C	C	C	C	C	AM	Wind direction	WNW	WNW	WNW	NW	NW
SSW	SW	W	WSW	SSE	NNW		PM						
Rainfall (mm) in last week				Rainfall (mm) from 01/01/2024 to till dated			Rainfall (mm) from 01/06/2024 to today						
70.3				2413.6			1782.8						

Weather summary / Alert: As per the forecast of IMD, there is a chance of moderate rainfall in isolated places of the district. Normal to above normal rainfall is expected in Kerala.

General and Agrometeorological Advisories for Thrissur District

Special crop management is required for agricultural crops during monsoon season. Proper drainage facilities have to be ensured. If the soil is more acidic, add lime as per soil testing. Pesticides should not be applied during rainy period. Provide supports to banana crops. Make the vegetable pandals stronger. To protect farmers and livestock from lightning and thunder avoid standing in open areas.

Crop	Crop/ Animal Advisory	Crop/ Animal Advisory
Rice (virippu) Panicle initiation stage	Bacterial Leaf blight	To control and prevent bacterial leaf blight in paddy, dissolve 20 g of cow dung in one litre of water and spray the supernatant liquid by mixing with 20g of pseudomonas.
Banana	Rhizome rot	During rainy season, there is a chance of Rhizome rot in banana. As a prophylactic measure, drench the basin with 3 litre of pseudomonas solution (20g per one liter of water).
Ginger	Phyllosticta leaf spot	Remove and destroying disease affected plants. As a prophylactic measure, spray 2ml hexaconazole or 1 ml Propiconazole or 2 gram SAAF or 2.5 g copper oxychloride or 2 g copper hydroxide dissolved in one litre of water. Otherwise 1% Bordeaux mixture can be sprayed. Spraying can be continued at fortnightly intervals depending upon the intensity of the disease.
Nutmeg	Leaf fall	During rainy season, there is a chance of leaf fall in nutmeg. As a prophylactic measure to manage leaf fall in nutmeg, spray 1% Bordeaux mixture. Proper drainage facilities have to be ensured

Animal Husbandry: Precautions should be taken to avoid ectoparasitic infestations. Deworming of animals should be done after microscopical examination of dung sample in a veterinary hospital. (CAADECCS, KVASU, Mannuthy)