


Advisory Service bulletin 34/16, Friday dt.19-08-2016

(Issued jointly by IAAS Vellanikkara)





THRISSUR

CURRENT SYNOPTIC SITUATION: The deep depression moved West wards and lies over south UttarPradesh and adjoining northeast Madhyapradesh near latitude 25⁰ N and longitude 82.3⁰ E. The feeble trough of low pressure is extending from Karnataka coast to Kerala coast..

NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST ISSUED ON: 19.08.2016 VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE: KERALA

PARAMETERS	ENSEMBLE FCST				
	DAY-1 20/08	DAY-2 21/08	DAY-3 22/08	DAY-4 23/08	DAY-5 24/08
DISTRICT : THRISSUR					
Rainfall (mm)	4	4	2	8	16
Max Temperature (deg C)	31	31	31	31	31
Min Temperature (deg C)	23	23	23	23	23
Total cloud cover (octa)	6	5	5	5	6
Max Relative Humidity (%)	90	90	90	90	90
Min Relative Humidity (%)	70	70	70	70	70
Wind speed (kmph)	800	800	800	800	008
Wind direction (deg)	290	320	320	290	290

Agro Meteorological Advisories

Crop	Stage of the crop / Pest/Disease	Advisories
		There is a chance for Sheath blight disease during rainy season.
Paddy	Panicle	The immediate symptom is that outer leaves of paddy become
(Jyothi,	initiation/	yellow coloured. Black lesions can be seen on the lower part
Kanchana)	Sheath blight	just above the water level. Apply Propiconazole 25 EC @ 500
		ml per 1 hector to control Sheath blight.
		There is a chance of yellowing in banana. Examine the sucker
Banana	Yellowing	and make sure that it is Erwinia rot, it can be controlled by
	(Erwinia Rot)	mixing 25gms of Kocide and 6gms of Streptocycline in 25 ml
		of water and apply 5 litre of solution to each banana basin.
		During rainy season there is a chance of rhizome rot, to control
Ginger, Turmeric	Rhizome rot	this dig out the affected plants and apply Kocide 2gm/1 litre
		water.
	D '41	To control fruit borer Apply any pesticide contains Neem oil (4
Cowpea	Fruit borer	ml per one liter of water).Repeat it in every week.