




Advisory Service bulletin 39/16, Friday dt.23-09-2016

(Issued jointly by IAAS Vellanikkara)

Dept. of Agriculture Meteorology, College of Horticulture, Vellanikkara

E mail: kauagmet@yahoo.co.in, cohagmet@kau.in



THRISSUR

CURRENT SYNOPTIC SITUATION: A well marked low pressure area over coastal Andhra Pradesh and neighbourhood lies over coastal Andhra Pradesh and adjoining areas of South Chattisgarh and Telangana. Associated upper air cyclonic circulation extends upto 7.6 km a.s.l. tilting southwards with height .

NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST

ISSUED ON: 23.09.2016

VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE : KERALA

PARAMETERS	ENSEMBLE FCST				
	DAY-1 24/09	DAY-2 25/09	DAY-3 26/09	DAY-4 27/09	DAY-5 28/09
DISTRICT : THRISSUR					
Rainfall (mm)	2	2	4	4	6
Max Temperature (deg C)	31	31	31	31	31
Min Temperature (deg C)	24	24	24	24	24
Total cloud cover (octa)	6	6	6	6	6
Max Relative Humidity (%)	90	90	90	90	90
Min Relative Humidity (%)	70	70	70	70	70
Wind speed (kmph)	013	013	008	008	008
Wind direction (deg)	290	290	270	270	270

Agro Meteorological Advisories:

Crop	Stage of the crop / Pest/Disease	Advisories
Paddy (Jyothi, Kanjana)	Harvesting stage/Rice bug	Apply neem based pesticides (3-4 ml per one liter of water). Pesticides application should be done either before 9 am or after 3 pm. Otherwise, the pesticide application will adversely affect the pollination.
Banana	Yellowing (Erwinia Rot)	There is a chance of yellowing in banana. Examine the sucker and make sure that it is Erwinia rot, it can be controlled by mixing 25gms of Kocide and 6gms of Streptocycline in 25 litre of water and apply 5 litre of solution to each banana basin.
Nutmeg	Leaf fall	Remove the infected leafs from field and apply Kocide 2g per one liter of water.
Cowpea	Anthraco nose	Apply Bavistin 1g per one liter of water.