







(Valid from 25.09.2024 to 29.09.2024)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 39/2024, (77) Tuesday

Dated 24.09.2024

			during las 24 th Sept		24)		Weather Parameters		Weather Forecast Valid for 25 ^{8h} September to 29 th September 2024				
18/09	19/09	20/09	21/09	22/09	23/09	24/09	Date		25/09	26/09	27/09	28/09	29/09
0.0	0.0	0.0	0.0	0.0	2.6	0.6	Rainfall (mm)		8	2	1	2	0.5
31.9	32.7	32.2	31.8	32.5	32.4	31.9	$T_{max}(^{0}C)$		33	33	33	33	33
24.4	23.8	23.6	23.6	23.5	24.4	23.4	T min(⁰ C)		23	23	25	25	25
Partly cloudy	Cloudy	Cloudy	Partly cloudy	Cloudy	Partly cloudy	Overcast	AM	Cloud	Clanda	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy
Partly cloudy	Partly cloudy	Partly cloudy	Cloudy	Cloudy	Cloudy	Cloudy	PM	cover	Cloudy				
89	93	97	90	93	90	95	RH-I (%)		86	86	86	86	86
60	59	68	62	61	81	73	RH-II (%)		68	68	68	68	68
2.2	1.9	1.8	1.6	1.6	2.1	2.6	Wind Speed (km/hr.)		6	6	5	4	4
Calm	Calm	Calm	Calm	Calm	Calm	NNE	AM	Wind	wsw	WSW	WSW	W	W
WNW	SW	WNW	NW	SW	Calm	ENE	PM	direction					
Ra	Rainfall (mm) in last week Rainfall (mm) from 0					Rainfall (mm) from 01/06/2024 to today							
3.2				2734.3					2103.5				

Weather summary / Alert: As per the forecast of IMD, there is a chance of light rainfall in the district. Above normal rainfall is expected in Kerala.

General and Agrometeorological Advisories for Thrissur District

Special crop management is required for agricultural crops during monsoon season. Proper drainage facilities have to be ensured. If the soil is more acidic, add lime as per soil testing. Pesticides should not be applied during rainy period. Provide supports to banana crops. Make the vegetable pandals stronger.

Crop	Pest/ Disease, Crop Management	Agromet Advisory					
Rice (<i>Virippu</i>) Maturity stage	Rice bug	There is a chance of appearance of rice bug in paddy. To control spray fish jaggery extract (20 ml per litre of water) or spray chitin based <i>Pseudomonas fluorescens</i> 20 gram in one litre of water. If the attack is severe, spray Azadirachtin 10 ml or Malathion 2 ml per litre of water.					
Rice (Mundakan)	Land preparation	Apply lime during land preparation. Seed treatment with <i>pseudomonas</i> can be done before sowing. (10 gram of <i>pseudomonas</i> for one kg of paddy seeds).					
Banana	Sigatoka leaf spot disease	As a prophylactic measure to sigatoka leaf spot disease, spray 20g of <i>pseudomonas</i> in one litre of water. If sigatoka attack is seen, spray one milli litre hexaconazole or one milli litre propiconazole per one litre of water on both the surface of the leaf.					
Pepper	Pollu disease	During high humidity, there is a chance of pollu disease in pepper plants. Control the shade in the garden and spray with 1% Bordeaux mixture.					

Animal Husbandry: - If leg lameness or fever was observed in cattle, medical attention should be given immediately. Thoroughly wash udder of cow and hands of milking person before milking using clean water with an added disinfectant, such as 1:1000 potassium permanganate solution. (CAADECCS, KVASU, Mannuthy).