


Agromet Advisory Service bulletin 45/16, Tuesday dt 01-11-16

(Issued jointly by IAAS Vellanikkara)

Dept. of Agriculture Meteorology, College of Horticulture, Vellanikkara E mail: kauagmet@vahoo.co.in, cohagmet@kau.in

THRISSUR



Significant weather for the preceding week at Vellanikkara

Rainfall weekly total	18.7 mm
Maximum temperature	31.7-32.8 [°] C
Minimum temperature	21.4-22.9 [°] C
Max. RH	91-98%
Min. RH	60-78%
Wind speed	0.5 -1.1 km/hr
Sun shine hour	0.8-8.0 hr
Evaporation	1.5-2.9 mm

CURRENT SYNOPTIC SITUATION: The upper air cyclonic circulation lay over Interior Tamilnadu and adjoining areas of Kerala and South Interior Karnataka now lies over Southeast Arabian sea off Kerala coast and extends upto 0.9km asl. A low pressure area has formed over south Andaman sea and neighbourhoos and its upper air cyclonic circulation extends upto 7.6 km asl.

HEAVY RAINFALL WARNING : Heavy rain is most likely to occur at one or two places in Kerala on 2^{nd} November 2016

NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST ISSUED ON: 01.11.2016 VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE: KERALA

PARAMETERS	ENSEMBLE FCST				
	DAY-1 02/11	DAY-2 03/11	DAY-3 04/11	DAY-4 05/11	DAY-5 06/11
DISTRICT : THRISSUR					
 Rainfall (mm)	16	2	3	3	0
Max Temperature (deg C)	30	30	31	31	31
Min Temperature (deg C)	22	23	23	23	23
Total cloud cover (octa)	6	6	6	5	5
Max Relative Humidity (%)	90	90	90	90	90
Min Relative Humidity (%)	70	70	70	70	70
Wind speed (kmph)	006	006	004	004	004
Wind direction (deg)	020	220	290	290	270

Agro Meteorological Advisories:

Сгор	Stage of the crop / Pest/Disease	Advisories
Paddy		There is a chance of rice case worm in paddy field. To control this 25 kg sawdust mixed with 1 litre kerosene can be
(Jyothi, kanjana, Uma, Ponmani)	Seedling stage	spread after draining the field for 1 acre. Fields which cannot be drained can be sprayed with fame (2ml/10L) or Coragen
		(3ml/ 10L). Soil can be made to a good tilth in coconut farms to enable
Coconut	-	the rain water infiltration and proper soil air movement. Measures should be taken to control weeds and root grub.
Banana	Sigatoka	There is a chance of sigatoka in banana field. Apply Tilt 1 ml/1 litre of water.
Vegetables	Stuted growth	There is a chance of stunted growth in bitter gourd. As a preventive measure spray cows urine 50 ml/ 1 L water.