

## Agromet Advisory Service bulletin 45/17, Friday dt.10-11-2017

(Issued jointly by IAAS Vellanikkara)

Dept. of Agricultural Meteorology, College of Horticulture, Vellanikkara Email: kauagmet@yahoo.co.in, cohagmet@kau.in

## THRISSUR

**CURRENT SYNOPTIC SITUATION:** A trough of low pressure area lies over south west Bay of Bengal and adjoining Sri Lanka and Tamil Nadu coast with associated upper air cyclonic circulation extending upto 3.1 km above sea level.

## NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST ISSUED ON: 10.11.2017 VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

DISTRICT: THRISSUR ENSEMBLE FCST

PARAMETERS	11.11.2017	12.11.2017	13.11.2017	14.11.2017	15.11.2017
Rainfall (mm)	7	0	5	5	11
Max Temperature ( deg	33	33	33	33	33
Min Temperature ( deg C)	22	22	22	22	22
Total cloud cover (octa)	6	6	6	6	6
Max Relative Humidity (%)	90	90	90	90	90
Min Relative Humidity (%)	70	70	70	70	70
Wind speed (kmph)	6	6	6	6	6
Wind direction (deg)	90	90	90	90	90

**Agro Meteorological Advisories:** 

Crop	Crop stage / Pest/Disease	Advisories		
Paddy		During this special weather condition, there is a chance of occurrence of		
(Jyothi, kanjana)		leaf folder and stemborer in mundakan paddy (second crop). Two cc		
	Leaf folder,	trichograma cards, which is cut into 10 pieces, can be placed under leaf		
	Stem borer	in different corners of field to control them. If the pest attack is severe, apply flubendiamide (fame) @ 2ml per 10 litre of water.		
		There is a chance of occurrence of diseases like brown spot and blast in		
	Brown spot,	this weather condition. Spray nativo 4 g per 10 litre of water to control		
Blast				
		them.		
		Destroy the affected leafs along with the spodoptera. For severe attack		
Banana	Spodoptera	spray flubendiamide (fame) 2 ml per10 litre of water.		
Vegetables	Epilachna	Apply quinalphos 2 ml/1 litre of water for epilachna beetle attack in		
(Snake gourd)	beetle	vegetables		
Animal Husbandry	If respiratory distress, fever or lameness are noticed in animals, consult a veterinary			
	surgeon immediately. Mosquitoes and rat breeding near cattle shed premises should be			
	avoided. (CAADECS, KVASU, Mannuthy)			