



Agromet Advisory Bulletin for Thrissur District

(Valid from 08.11.2025 to 12.11.2025)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara
based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 45/2025, (89) Friday

Dated 07.11.2025

Medium range weather forecast of IMD for next five days

Weather parameters	08.11.2025	09.11.2025	10.11.2025	11.11.2025	12.11.2025
Rainfall (mm)	1	2	0.1	0.1	0.2
Maximum temperature (°C)	33	33	33	33	33
Minimum temperature (°C)	22	22	22	22	22
Maximum Relative Humidity (%)	75	75	75	75	75
Minimum Relative Humidity (%)	70	70	70	70	70
Wind speed (kmph)	3	3	2	3	3
Wind direction (deg)	270	290	200	200	270
Total cloud cover (octa)	8	8	7	8	8

Weather summary / Alert: As per the forecast of IMD, there is a chance of light rainfall in the district. Below normal rainfall is expected in Kerala. Thunderstorm accompanied by lightning is very likely to occur at one or two places in Kerala.

General and Agrometeorological Advisories for Thrissur District

General Advisory: -Prophylactic measures can be taken to control crop pest & diseases. Proper mulching of crops can be followed to conserve soil moisture.

Crop	Pest/ Disease, Crop Management	Agromet Advisory
Rice (Mundakan) Seedling stage	Fertilizer application	The first dose of fertilizer can be applied in paddy. For one acre of paddy field, 51.5 kg of urea, 77.7 kg of rock phosphate, and 15 kg of muriate of potash can be applied. (If Factamfos is used instead of rock phosphate, 21.18 kg of urea is sufficient.)
	Thrips	To control thrips in rice, spray 2 gram of Thiomethoxam in 10 litre of water.
Banana	Rhizome Weevil	There is a chance of Rhizome Weevil attack in banana. As a precaution, cut and remove the outer layer of the rhizome before planting and sundry for 3-4 days after smearing it with slurry of cow dung and ash. Apply neem cake @ 1 kg/plant at the time of planting.
Turmeric	Leaf Blotch (<i>Taphrina maculans</i>)	There is a chance of leaf blotch diseases in turmeric. Remove and destroy disease affected plants. As a prophylactic measure, spray 2ml hexaconazole or 2 gram SAAF dissolved in one litre of water. Spraying can be continued at fortnightly intervals depending upon the intensity of the disease.

Animal Husbandry: - Vaccinate animals against foot and mouth disease and hemorrhagic septicemia. If needed, give medicine against worm infestation after dung examination. (CAADECCS, KVASU, Mannuthy).