



Kerala Agricultural University, Vellanikkara, Thrissur
Department of Agricultural Meteorology, College of Horticulture

Email: cohagmet@kau.in&kauagmet@yahoo.co.in

Agromet Advisory Service Bulletin

(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)



THRISSUR DISTRICT

AAS 47/2020,(94) Friday

20.11.2020

SIGNIFICANT WEATHER FOR THE PRECEDING WEEK AT VELLANIKKARA:

The total rainfall received 8.8 mm. The maximum temperature ranges from 31.2°C to 34.4°C, minimum temperature ranges from 21.0°C to 24.0°C. Morning relative humidity ranges from 67% to 86% and afternoon relative humidity ranges from 53% to 69%. Wind speed ranges from 3.0 to 10.5 km/h. Bright sunshine hours range from 2.6 to 6.6 hours. Evaporation ranges from 3.1 to 5.2 mm.

CURRENT SYNOPTIC SITUATION: A low pressure area lies over central parts of south Arabian Sea with associated cyclonic circulation extending upto 4.5 km above mean sea level. A cyclonic circulation lies over Equatorial Indian Ocean and adjoining Southeast Bay of Bengal extending upto 3.1 km above mean sea level.

WEATHER FORECAST FOR THE NEXT FIVE DAYS:

Weather parameters	21.11.2020	22.11.2020	23.11.2020	24.11.2020	25.11.2020
Rainfall (mm)	7	0	0	0	0
Maximum temperature(°C)	33	33	33	33	33
Minimum temperature (°C)	22	22	22	22	22
Maximum Relative Humidity (%)	83	83	83	83	83
Minimum Relative Humidity (%)	60	60	60	60	60
Wind speed (kmph)	4	14	18	16	8
Wind direction (deg)	140	50	50	50	20
Total cloud cover (octa)	6	3	3	2	5

As per the forecast of IMD & KAU, there is a chance of less rainfall in the district.

AGRO-METEOROLOGICAL ADVISORIES:

Crop	Pest/ Disease	Advisories
General Advisories: precaution should be taken during drying of agricultural products and spraying of pesticides.		
Rice (Mundakan) Tillering stage	Bacterial Leaf blight	To control the spread of Bacterial Leaf blight in paddy, spray 6 g of Streptocycline in 30 litre of water.
	Iron toxicity	If there is toxicity of iron in the rice field, apply 150 kg lime per acre.
Banana	Spodoptera	There is a chance of attack of spodoptera in banana. Destroy the affected leaves along with the spodoptera.
Turmeric	Leaf Blotch	1 % Bordeaux mixture can be sprayed. Otherwise spray 2.5 g mancozeb or 1g carbendazim or 2.5 g copper oxychloride (Blitox or Blue copper) or 2ml hexaconazole (Contaf) or 0.75 ml difenconazole (Score) dissolved in one litre of water. Spraying can be continued at fortnightly intervals depending upon the intensity of the disease.
Vegetables (cowpea)	Fruit borer	To control fruit borer in cowpea, apply 5% Neem Kernel Suspension to control fruit borer in the initial stage. Spray Flubendamide 2 ml per10 litre of water or Chlorantraniliprole 3 ml per10 litre if the attack is severe.
Animal Husbandry	Animal should not be allowed to graze in open grazing lands to avoid lightening risk. Vaccinate animals against foot & mouth disease and haemorrhagic septicaemia	