




## Advisory Service bulletin 50/16, Friday dt.09-12-2016

( Issued jointly by IAAS Vellanikkara)

Dept. of Agriculture Meteorology, College of Horticulture, Vellanikkara

E mail: kauagmet@yahoo.co.in, cohagmet@kau.in

**THRISSUR**



**CURRENT SYNOPTIC SITUATION:** The cyclonic storm “VARDAH” lies over South east bay of Bengal and centered at 0830 hrs IST of today near latitude 12.1<sup>0</sup> N and longitude 90.4<sup>0</sup> E.

### NWP MODELS BASED DISTRICT LEVEL WEATHER FORECAST

ISSUED ON: 09.12.2016

VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE : KERALA

PARAMETERS	ENSEMBLE FCST				
	DAY-1 10/12	DAY-2 11/12	DAY-3 12/12	DAY-4 13/12	DAY-5 14/12
DISTRICT : THRISSUR					
Rainfall (mm)	0	0	0	0	0
Max Temperature ( deg C)	34	34	34	34	34
Min Temperature ( deg C)	23	23	23	22	22
Total cloud cover (octa)	5	5	4	4	4
Max Relative Humidity (%)	80	80	80	80	80
Min Relative Humidity (%)	60	60	60	60	60
Wind speed (kmph)	006	006	006	006	006
Wind direction (deg)	290	290	320	050	050

### Agro Meteorological Advisories:

Crop	Stage of the crop / Pest/Disease	Advisories
Paddy (Jyothi, kanjana, Uma, Ponmani)	Panicle initiation/Sheath blight, Stem borer	Since cloudy condition prevailed, there is a chance of Sheath blight in paddy field. Spray Tilt 1 ml/1 litre of water. To control Stem borer, spray Regent 2 gm/ 1 litre of water.
Banana	Pseudostem weevil	Apply 2% neem oil-garlic-soap emulsion in the starting stage. In case of severe attack apply 3ml Chlorpyrifos in 1 liter of water on the pseudostem mixing with gum.
Pepper	Quick wilt	There is a chance of Quick wilt disease. Spray 20g pseudomonas/ 1 litre of water.
Cowpea	Fruit borer	To control fruit borer Apply any pesticide contains Neem oil (4 ml per one litre of water).Repeat it in every week.