



Kerala Agricultural University, Vellanikkara, Thrissur
Department of Agricultural Meteorology, College of Horticulture

Email: cohagmet@kau.in&kauagmet@yahoo.co.in

Agromet Advisory Service Bulletin

(Issued jointly by AMFU Thrissur, India Meteorological Department and Department of Agriculture)



THRISSUR DISTRICT

AAS 50/2020, (100) Friday

11.12.2020

SIGNIFICANT WEATHER FOR THE PRECEDING WEEK AT VELLANIKKARA:

The total rainfall received 7.7 mm. The maximum temperature ranges from 27.9°C to 32.7°C, minimum temperature ranges from 21.0°C to 23.3°C. Morning relative humidity ranges from 71% to 93% and afternoon relative humidity ranges from 47% to 70%. Wind speed ranges from 1.8 to 10.2 km/h. Bright sunshine hours range from 0.0 to 8.1 hours. Evaporation ranges from 1.2 to 4.5 mm.

CURRENT SYNOPTIC SITUATION: A low pressure area lies over central parts of south Arabian sea with associated cyclonic circulation extending upto 3.1 km above mean sea level.

WEATHER FORECAST FOR THE NEXT FIVE DAYS:

Weather parameters	12.12.2020	13.12.2020	14.12.2020	15.12.2020	16.12.2020
Rainfall (mm)	0	0	0	0	0
Maximum temperature(°C)	33	33	33	33	33
Minimum temperature (°C)	22	22	22	22	22
Maximum Relative Humidity (%)	74	74	74	74	74
Minimum Relative Humidity (%)	53	53	53	53	53
Wind speed (kmph)	14	10	4	8	14
Wind direction (deg)	50	50	20	50	50
Total cloud cover (octa)	4	2	1	1	1

As per the forecast of IMD & KAU, no rainfall is forecasted in the district.

AGRO-METEOROLOGICAL ADVISORIES:

Crop	Pest/ Disease	Advisories
General Advisories: Since dry weather condition is prevailed, irrigation to crops should be ensured. Drying of agricultural products and spraying of pesticides can be done.		
Rice (Mundakan) Panicle Emergence	Bacterial Leaf blight	As a prophylactic measure, spray 20 gram Pseudomonas in one litre of water. To control the spread of Bacterial Leaf blight in paddy, spray 6 g of Streptocycline in 30 litre of water.
	Leaf folder	There is a chance of leaf folder attack in mundakan Paddy. It can be controlled by using trichogramma cards (2CC per one acre). Cut the cards in to small pieces and fix in different regions of the field. If the attack is severe, spray Chlorantraniliprole (Coragen) @ 3ml per ten litre of water.
	Brown Plant Hopper (BPH)	There is a chance of Brown plant hopper attack in rice. Spray 2 g Thiamethoxam (Actara) per ten litre of water.
Banana	Spodoptera	If the attack is in the initial stage, the affected leaves should be destroyed along with egg, caterpillar and pupa. Neem Kernel suspension (5 percentage) should be sprayed.
Vegetables (cowpea)	Fruit borer	If the attack is in the initial stage, the affected leaves should be destroyed along with egg, caterpillar and pupa. Neem Kernel suspension (5 percentage) should be sprayed. If the attack is severe, spray Flubendamide (Tukumi) 2 ml per ten litre of water or Chlorantraniliprole (Coragen) 3 ml per ten litre of water.
Cabbage/ Cauliflower	Spodoptera	
Animal Husbandry	1. Take precautions to avoid ecto parasitic infestation. 2. If fever, shivering, brown coloured urine, off feed etc. are noticed, consult veterinary doctor immediately.	