



Agromet Advisory Bulletin for Thrissur District

(Valid from 21.12.2022 to 25.12.2022)

Prepared by the Department of Agricultural Meteorology, Kerala Agricultural University, COA, Vellanikkara
based on the medium range weather forecast of India Meteorological Department (IMD), New Delhi

AAS Bulletin No: 51/2022, (101) Tuesday

dated 20.12.2022

A. Significant weather for the preceding week at vellanikkara

Rainfall (mm)	Maximum Temp.(°C)	Minimum Temp.(°C)	RH FN (%)	RH AN (%)	Wind speed (Km/h)	Sun shine hours	Evap. (mm)
(3.8)	(29.2-33.3)	(21.1-24.0)	68-90	(38-72)	(0.6-5.1)	(1.0-9.8)	(1.5-5.2)

B. Medium range weather forecast of IMD for next five days

Weather parameters	21.12.2022	22.12.2022	23.12.2022	24.12.2022	25.12.2022
Rainfall (mm)	0	0	0	0	0
Maximum temperature (°C)	32	32	32	32	32
Minimum temperature (°C)	24	24	24	22	22
Maximum Relative Humidity (%)	80	80	80	80	80
Minimum Relative Humidity (%)	65	65	65	65	65
Wind speed (kmph)	6	3	2	3	3
Wind direction (deg)	290	290	290	250	320
Total cloud cover (octa)	2	2	2	2	2

Weather summary / Alert: Above normal rainfall is expected in Kerala.

RAINFALL FORECAST FOR NEXT FIVE DAYS

2022 December 21	2022 December 22	2022 December 23	2022 December 24	2022 December 25
No rain	No rain	No rain	No rain	No rain

C. General and Agrometeorological Advisories for Thrissur District

Prophylactic measures to control crop pest & diseases can be taken to avoid their spread. Since there is a chance of dry weather condition due to strong wind and high evaporation, crops should be irrigated, when required. Since the wind speed may increase, give strong propping to banana and staking to vegetables.

Crop	Pest/ Disease	Crop/ Animal Advisory
Rice (Mundakan) Panicle emergence stage	Sheath Blight	To control sheath blight disease in rice, spray 4 gram of (Trifloxystrobin + Tebuconazole) mixture in ten litres of water.
Banana	Spodoptera	If the attack is in the initial stage, the affected leaves should be destroyed along with egg, caterpillar and pupa. Neem Kernel suspension (5 percentage) should be sprayed.
Cabbage/ Cauliflower	Spodoptera	If the attack is in the initial stage, the affected leaves should be destroyed along with egg, caterpillar and pupa. Neem Kernel suspension (5 percentage) should be sprayed. If the attack is severe, spray Flubendamide 2 ml per ten litre of water or Chlorantraniliprole 3 ml per ten litre of water.

Animal Husbandry- Care should be taken to avoid ecto-parasitic infestation. Ensure sufficient supply of clean drinking water to the animals. (CAAECCS, KVASU, Mannuthy).